GREENBELT SValagro

CRIENE BEIN

NET WEIGHT 5Lb (2.3 Kg)

GliMO Fe 0-0-0

Iron (Fe)	6.0%
6.0% Chelated Iron (Fe)	
Manganese(Mn)	1.0%
1.0% Chelated Manganese (Mn)	
Derived from Iron EDDHA: Manganese EDTA	

GENERAL INFORMATION

GREENBELT (0-0-0) is a source of Iron and Manganese chelates readily available for crops, especially in alkaline soil conditions. The product is to be applied into the root zone through drip line, drench, injection, band application (watering in the solution).

Produced and packed by VALAGRO S.p.A.
Zona Ind.le - 66041 Atessa (Ch) - ITALY
Distributed and guaranteed by Valagro USA Inc.
19500 Hwy 249, suite 245, Houston TX 77070
Tel (281) 664 8700 Fax (281) 664 8701

GREENBELT is a Registered Trademark of VALAGRO S.p.A.
Produced and packed by VALAGRO S.p.A.
Zona Industriale - 66041 Atessa (Ch) - ITALY
Distributed and guaranteed by Valagro USA Inc.
19500 Hwy 249, suite 245, Houston TX 77070
Tel (281) 664 8700 Fax (281) 664 8701





Ш

œ



GREENBELT

F1873 ????????????? 00

NOTICE: Direct exposure to sunlight is to be avoided.

KEEP OUT OF REACH OF CHILDREN

CONDITIONS OF SALE

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with direction under normal condition and use. No one other than an office of the seller is authorized to make any warranty, guaranty or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price. It is impossible to eliminate all risks associated with the use of this product. Crop injury, lack of performance, all other unintended consequences may result because of such factors as use of the product contrary to label instruction, abnormal condition, presence of other material, the manner of application, or other factors, all of which are beyond the control of Valagro USA or the seller. All such risks shall be assumed by buyer.

STORAGE: Keep the product sheltered from the sun and in a dry warehouse. When applied do not leave exposed to sunlight and cover immediately with soil. Once open, completely use the contents of the bag.

FIELD APPLICATION INSTRUCTIONS

Note: The dosage range indicated is dependent upon the plant size and the severity of chlorosis.

CROP	RATE	TIMING
Fruit crops	5 - 15 lbs./acre per application	2-3 treatments starting from vegetative growth
Industrial Crops	5 - 15 lbs./acre per application	At planting or when deficiency symptoms first appear
Vegetable Crops	5 - 15 lbs./acre per application	1-2 treatments according plant needs after planting
Floriculture	4-6 oz / 1000 sq.ft.	At planting or when deficiency symptoms first appear

→ Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.htm







